

MICROBIOME, METABOLOMICS, METAGENOMICS

Dietary gluten-induced gut dysbiosis is accompanied by selective upregulation of microRNAs with intestinal tight junction and bacteria-binding motifs in rhesus macaque model of celiac disease.

Mohan M., Chow C.-E.T., Ryan C.N., Chan L.S., Dufour J., Aye P.P., Blanchard J., Moehs C.P., Sestak K.

Nutrients. 2016 Oct 28;8(11).

<https://www.ncbi.nlm.nih.gov/pubmed/27801835>

Salivary and fecal microbiota and metabolome of celiac children under gluten-free diet.

De Angelis M., Vannini L., Di Cagno R., Cavallo N., Minervini F., Francavilla R., Ercolini D., Gobbetti M.

International Journal of Food Microbiology. 2016 Dec 19;239:125-132.

<https://www.ncbi.nlm.nih.gov/pubmed/27452636>

Lack of evidence of rotavirus-dependent molecular mimicry as a trigger of coeliac disease.

Ziberna F., De Lorenzo G., Schiavon V., Arnoldi F., Quaglia S., De Leo L., Vatta S., Martelossi S., Burrone O.R., Ventura A., Not T.

Clinical and Experimental Immunology. 2016 Dec;186(3):356-363.

<https://www.ncbi.nlm.nih.gov/pubmed/27548641>

Gut Microbiota and Risk of Developing Celiac Disease.

Cenit M.C., Codoner-Franch P., Sanz Y.

Journal of Clinical Gastroenterology. 2016 Nov/Dec;50 Suppl 2:S145-S147.

<https://www.ncbi.nlm.nih.gov/pubmed/27741161>

The Interaction between Celiac Disease and Intestinal Microbiota.

Losurdo G., Principi M., Iannone A., Ierardi E., Di Leo A.

Journal of Clinical Gastroenterology. 2016 Nov/Dec;50 Suppl 2:S145-S147.

<https://www.ncbi.nlm.nih.gov/pubmed/27741161>

The industrial food additive, microbial transglutaminase, mimics tissue transglutaminase and is immunogenic in celiac disease patients.

Matthias T., Jeremias P., Neidhofer S., Lerner A.

Autoimmunity Reviews. 2016 Dec;15(12):1111-1119.

<https://www.ncbi.nlm.nih.gov/pubmed/27640315>

No change in the mucosal gut microbiome is associated with celiac disease-specific microbiome alteration in adult patients.

D'Argenio V., Casaburi G., Precone V., Pagliuca C., Colicchio R., Sarnataro D., Discepolo V., Kim S.M., Russo I., Del Vecchio Blanco G., Horner D.S., Chiara M., Pesole G., Salvatore P., Monteleone G., Ciacci C., Caporaso G.J., Jabri B., Salvatore F., Sacchetti L.

Am J Gastroenterol. 2016 Nov;111(11):1659-1661

<https://www.ncbi.nlm.nih.gov/pubmed/27808136>

Changes in duodenal tissue-associated microbiota following hookworm infection and consecutive gluten challenges in humans with coeliac disease.

Giacomin P, Zakrzewski M, Jenkins TP, Su X, Al-Hallaf R, Croese J, de Vries S, Grant A, Mitreva M, Loukas A, Krause L, Cantacessi C.

Sci Rep. 2016 Nov 9;6:36797.

<https://www.ncbi.nlm.nih.gov/pubmed/27827438>